Bio-traps and Site Assessment Strategies for Groundwater Impacted by Chlorinated Hydrocarbons

> Kerry Sublette University of Tulsa

Dora Taggart, Brett Baldwin, Anita Biernacki, Kate Clark

Microbial Insights, Inc





What Are Bio-Trap[®] Samplers?

Passive sampling tool for microbes

Collects active microbes

Integrated sample vs. "snapshot"

Analyzed using molecular biological tools, analytical chemistry, and stable isotope analysis





How Do Bio-Traps Work?

Properties of Bio-Sep Beads

- 3-4 mm in diameter
- 25% Nomex and 75% PAC
- 74% porosity
- 600 m² of surface area/g
- Heat sterilized 270 °C
- Colonized by active microbes







Types of Bio-Trap Samplers

Standard Bio-Trap

- Basic design
- Sampling groundwater, surface
 - waters, sediments, soils
- Compatible with all MBTs, analytical chemistry, and stable
 - isotope techniques

















Amendments Include:

Electron Donors

- Vegetable oil
- Molasses
- HRC
- EOS
- Lactate
- And more

Electron Acceptors

- Oxygen (PermeOx, ORC)
- Nitrate
- Iron (III)
- Sulfate
- And more

Stable Isotope Compounds (¹³C)

- Benzene
- Toluene
- p-Xylene
- MTBE
- TBA
- Naphthalene
- Chlorobenzene
- 1,4-Dioxane
- Sulfolane
- And more





How Are Bio-Trap Samplers Deployed?

Purge monitoring well

Suspend from top of casing

Deploy within the screened interval at depths of interest.

If large fluctuations in the water level are anticipated suspended from a float.

Monitoring Well





How Are Bio-Trap Samplers Analyzed:

Molecular Biological Tools

- PLFA
- CENSUS (qPCR)
- QuantArrays
- DGGE
- Stable Isotope Probing (SIP)

Chemical Analysis

- Compound specific isotope analysis (CSIA)
- Dissolved Inorganic Carbon (DIC)
- Contaminant Concentrations





What Can I Do With a Bio-Trap Sampler?

- Determine if known degraders of a COC are present
- Evaluate monitored natural attenuation versus enhanced bioremediation
- Compare effectiveness of amendments designed to stimulate bioremediation
- Prove that bioremediation of a specific compound is occurring



What Can I Do With a Bio-Trap Sampler?

- Determine if known degraders of a COC are present
- Evaluate monitored natural attenuation versus enhanced bioremediation
- Compare effectiveness of amendments designed to stimulate bioremediation
- Prove that bioremediation of a specific compound is occurring



Microbial Insights Database - Dehalococcoides





Determine If Known Degraders Are Present



CENSUS[®] - *Dehalococcoides* populations indicate the potential for complete reductive dechlorination of PCE to ethene but stimulation needed



What can I do with a Bio-Trap Sampler?

- Determine if known degraders of a COC are present
- Evaluate monitored natural attenuation versus enhanced bioremediation
- Compare effectiveness of amendments designed to stimulate bioremediation
- Prove that bioremediation of a specific compound is occurring











Screening Remediation Options: Chlorinated hydrocarbon impacted site

Control (MNA)

BioStim

(Electron

Donor)

Control Unit

- Evaluate MNA as treatment alternative
- Baseline for enhanced remediation options

BioStim Unit – Electron Donor Addition

- Enhanced anaerobic bioremediation
- Lactate, HRC, EOS

BioAug Unit – Culture and Electron Donor

Bioaugmentation (culture impregnated in beads in the bio-trap)



BioAug (Electron donor + culture)



Case Study: Amended ISMs



Chlorinated Solvent Site



Site Background

- Shallow aquifer impacted by chlorinated solvents, primarily trichloroethene (TCE).
- Daughter product *cis*-1,2 dichloroethene (DCE) has been detected.
- DCE appears to be accumulating with no observed production of vinyl chloride or ethene ("DCE stall").
- Biostimulation (electron donor addition) and bioaugmentation (donor and culture) were being considered as remediation strategies.



Site Specific Questions

Microbiology

Are organisms capable of complete reductive dechlorination of TCE to ethene (*Dehalococcoides*) present under MNA conditions?

Will addition of an electron donor stimulate growth of these key dechlorinating bacteria?

Will a bioaugmentation culture survive? Is bioaugmentation necessary?



Site Specific Questions

Chemistry

Will electron donor addition promote daughter product formation and stimulate complete reductive dechlorination?

Will bioaugmentation + biostimulation more effectively stimulate reductive dechlorination than biostimulation alone?







Control (MNA) Unit – CENSUS® qPCR Results





MNA vs BioStim





MNA vs BioStim vs BioAug





Control vs. BioStim – Impact on COCs









Bio-traps and Chlorinated Hydrocarbon Impacted Sites

- Standard bio-traps can measure degradation potential by quantifying *Dehalococcoides* and functional genes characteristic of reductive dechlorination (pre- or postinjection)
- Bio-traps coupled with *in situ* microcosms can compare effectiveness of amendments designed to stimulate bioremediation

For a copy of this presentation email <u>kerry-</u> <u>sublette@utulsa.edu</u>

For more information go to microbe.com

